

Single production of the doubly charged *Higgs* boson via $e\gamma$ collision in the *Higgs* triplet model

Chong-Xing Yue, Xue-Song Su, Jiao Zhang, Jue Wang

Department of Physics, Liaoning Normal University, Dalian 116029, P. R. China *

October 25, 2010

Abstract

The *Higgs* triplet model (*HTM*) predicts the existence of a pair of doubly charged *Higgs* bosons $H^{\pm\pm}$. Single production of $H^{\pm\pm}$ via $e\gamma$ collision at the next generation e^+e^- International Linear Collider (*ILC*) and the Large Hadron electron Collider (*LHeC*) is considered. The numerical results show that the production cross sections are very sensitive to the neutrino oscillation parameters. Their values for the inverted hierarchy mass spectrum are larger than those for the normal hierarchy mass spectrum at these two kinds of collider experiments. With reasonable values of the relevant free parameters, the possible signals of the doubly charged *Higgs* bosons predicted by the *HTM* might be detected in future *ILC* experiments.

PACS number: 12.60.Cn, 14.80.Cp, 13.10.+q

*E-mail: cxyue@lnnu.edu.cn

1. Introduction

During the past decade, neutrino oscillation experiments have provided us with very convincing evidence that neutrinos are massive particles mixing with each other [1]. This exciting breakthrough gives a strong motivation for physics beyond the standard model (SM), since the SM itself only contains three massless neutrinos [2]. Various ways to go beyond the SM have been proposed in order to accomodate this observation. The *Higgs* triplet model (HTM) [3, 4] is a simple extension of the SM with an $SU(2)_L$ -triplet *Higgs* boson of hypercharge $Y = 2$ whose vacuum expectation value (VEV) ν_Δ provides Majorana neutrino masses without introducing right-handed neutrinos.

The HTM [3, 4] generates a Majorana neutrino mass via the product of a triplet *Yukawa* coupling h_{ij} and a triplet VEV ν_Δ . This model predicts the existence of seven physical *Higgs* bosons which are two CP -even neutral bosons h^0 and H^0 , a CP -odd neutral one A^0 , a pair of single charged bosons $H^\pm$, and a pair of doubly charged *Higgs* bosons $H^{\pm\pm}$. These new particles can produce rich phenomenology at present and in future high energy collider experiments. Since charge conservation prevents doubly charged scalars from decaying to a pair of quarks, $H^{\pm\pm}$ carry the lepton number violation and would give rise to a distinctive experimental signal, which has lower background. This fact has lead to many studies involving the doubly charged *Higgs* bosons $H^{\pm\pm}$ in the literature. For example, it has been shown that such particles can be produced with sizable rates at hadron colliders via the processes $q\bar{q} \rightarrow H^{++}H^{--}$ [5, 6] and $q\bar{q}' \rightarrow H^{\pm\pm}H^\mp$ [5, 7, 8]. Using these production mechanisms, discovery of $H^{\pm\pm}$ at the Large Hadron Collider (LHC) has been studied in Refs. [9, 10].

There is a direct connection between the *Yukawa* coupling h_{ij} and the neutrino mass matrix which gives rise to phenomenological prediction for processes which depend on h_{ij} [11]. The branching ratios of the lepton flavor violating (LFV) decay processes $l_i \rightarrow l_j l_l l_k$ and $l_i \rightarrow l_j \gamma$ contributed by the doubly charged *Higgs* bosons $H^{\pm\pm}$ are strong depend on h_{ij} in the context of the HTM . Recently, it has been shown [12] that the current upper experimental limits for some of these LFV decay processes can give serve constraints on the parameter space of h_{ij} in the HTM . Taking into account these constraints, in

this paper, we consider single production of $H^{\pm\pm}$ via $e\gamma$ collision at the next generation e^+e^- International Linear Collider (*ILC*) [13, 14] and the Large Hadron electron Collider (*LHeC*) [15]. Although single production of the doubly charged *Higgs* bosons via $e\gamma$ collision has been studied in Refs. [16], in this paper, we will focus our attention on the effects of the neutrino oscillation parameters on its production cross section and see whether the neutrino mass hierarchy can be disentangled via this kind of production processes at the *ILC* and *LHeC*. Our numerical results show that the production cross sections at these two kinds of high energy colliders are very sensitive to the neutrino oscillation parameters. The production cross section of the process $e\gamma \rightarrow e^+H^{--}$ for the inverted hierarchy mass spectrum is significantly larger than that for the normal hierarchy mass spectrum. In the sizable range of the parameter space of the *HTM*, their values can be large enough to be detected in the future *ILC* experiments.

Our paper is organized as follows. The essential features of the *HTM* are briefly reviewed in section 2. In this section, we also summarized the constraints on the *HTM* from some *LFV* decay processes. The effective production cross sections for the subprocess $e\gamma \rightarrow l^+H^{--}$ at the *ILC* and *LHeC* are calculated in sections 3 and 4. Our conclusions and discussions are given in section 5.

2. The essential features of the HTM

The *Higgs* triplet model (*HTM*) [3, 4] is one of the appealing scenarios for generating neutrino masses without the introduction of a right-handed neutrino. In this model, a complex $SU(2)_L$ triplet scalar with the hypercharge $Y = 2$ is introduced to the *SM*, which can be parameterized by

$$\Delta = \begin{pmatrix} \Delta^+/\sqrt{2} & \Delta^{++} \\ \Delta^0 & -\Delta^+/\sqrt{2} \end{pmatrix}. \quad (1)$$

A nonzero *VEV* value $\langle \Delta^0 \rangle$ gives rise to the following mass matrix for neutrinos

$$m_{ij} = 2h_{ij} \langle \Delta^0 \rangle = \sqrt{2}h_{ij}\nu_\Delta. \quad (2)$$

The symmetric complex matrix h_{ij} ($i, j = e, \mu, \tau$) is the *Yukawa* coupling strength. The

interaction of the triplet scalar with lepton doublet $L_i \equiv (\nu_{iL}, l_{iL})^T$ is given by

$$\mathcal{L} = -h_{ij}[\nu_{iL}^T C \nu_{jL} \Delta^0 - \frac{1}{\sqrt{2}}(\nu_{iL}^T C l_{jL} + l_{iL}^T C \nu_{jL}) \Delta^+ - l_{iL}^T C l_{jL} \Delta^{++}] + h.c., \quad (3)$$

here C is the charge conjugation operator.

The neutrino mass matrix m_{ij} can be diagonalized by the Maki-Nakagawa-Sakata (MNS) matrix V_{MNS} [17]. Then the couplings h_{ij} can be written as following

$$h_{ij} = \frac{m_{ij}}{\sqrt{2}\nu_\Delta} \equiv \frac{1}{\sqrt{2}\nu_\Delta} [V_{MNS} \text{diag}(m_1, m_2 e^{i\varphi_1}, m_3 e^{i\varphi_2}) V_{MNS}^T]_{ij} \quad (4)$$

with

$$V_{MNS} = \begin{pmatrix} c_{12}c_{13} & c_{13}s_{12} & e^{-i\delta}s_{13} \\ -c_{12}s_{13}s_{23}e^{i\delta} - c_{23}s_{12} & c_{12}c_{23} - e^{i\delta}s_{12}s_{13}s_{23} & c_{13}s_{23} \\ s_{12}s_{23} - e^{i\delta}c_{12}c_{23}s_{13} & -c_{23}s_{12}s_{13}e^{i\delta} - c_{12}s_{23} & c_{13}c_{23} \end{pmatrix} \times \text{diag}(e^{i\varphi_1/2}, 1, e^{i\varphi_2/2}). \quad (5)$$

Where m_1, m_2, m_3 are the absolute masses of the three neutrinos. The phase δ is the Dirac CP-violating phase, φ_1 and φ_2 are referred to as the Majorana phases [4, 18] and there are $0 \leq \varphi_1, \varphi_2 < 2\pi$. $s_{ij} = \sin \theta_{ij}$, $c_{ij} = \cos \theta_{ij}$ with $0 \leq \theta_{ij} \leq \pi/2$, in which θ_{ij} are the mixing angles. The explicit expression forms for h_{ij} have been given in Refs. [8, 9].

After the eletroweak symmetry breaking, there are seven physical massive *Higgs* bosons ($H^{\pm\pm}, H^\pm, h^0, H^0, A^0$) left in the spectrum for the *HTM*. The doubly charged *Higgs* bosons $H^{\pm\pm}$ are entirely composed of the triplet scalars $\Delta^{\pm\pm}$, while the remaining eigen states are mixtures of the doublet and triplet scalars. Such mixing is proportional to the triplet *VEV* ν_Δ , which is generally small even if ν_Δ assumes its largest value of a few GeV. Thus, $H^\pm, H^0, A^0$ are predominantly composed of the triplet scalars, while the *SM*-like *Higgs* boson h^0 is mainly composed of the doublet scalars.

It is well known that the doubly charged *Higgs* bosons $H^{\pm\pm}$ have contributions to the *LFV* decays $l_i \rightarrow l_j \gamma$, $l_j \rightarrow l_j l_k l_l$ [19]. In the *HTM*, their branching ratios depend on the neutrino mass matrix parameters, the *VEV* value ν_Δ , and the mass $m_{H^{\pm\pm}}$ of the doubly charged Higgs bosons [12]. The current experimental upper limits for some of these *LFV*

decay processes can give severe constraints on the relevant free parameters. In the *HTM*, the *LFV* decay process $\mu \rightarrow e\gamma$ proceed via exchange of $H^{\pm\pm}$ and $H^\pm$ at one-loop. In the case of ignoring the electron mass in the final state and all lepton masses in the loop, the branching ratio $Br(\mu \rightarrow e\gamma)$ can be written as [12]

$$Br(\mu \rightarrow e\gamma) \approx \frac{27\alpha |(h^+h)_{e\mu}|^2}{64\pi G_F^2 m_H^4}, \quad (6)$$

where $\alpha = 1/137$ is the fine structure constant, $G_F = 1.17 \times 10^{-5} GeV^{-2}$ is the Fermi constant. In above equation, it has been assumed $m_{H^{\pm\pm}} = m_{H^\pm} = m_H$. The current experimental upper limit for the *LFV* process $\mu \rightarrow e\gamma$ is $Br^{exp}(\mu \rightarrow e\gamma) \leq 1.2 \times 10^{-11}$ [20], then we have

$$|(h^+h)_{e\mu}|^2 \leq 2.6 \times 10^{-9} \left(\frac{m_H}{200 GeV}\right)^4. \quad (7)$$

Using Eq.(4), there is

$$2\nu_\Delta^2 (hh^+)_{ij} = m_1^2 \delta_{ij} + [V_{MNS} diag(0, \Delta m_{21}^2, \Delta m_{31}^2) V_{MNS}^+]_{ij}, \quad (8)$$

where $\Delta m_{21}^2 = m_2^2 - m_1^2$ and $\Delta m_{31}^2 = m_3^2 - m_1^2$. Since the sign of Δm_{31}^2 is undetermined at present, distinct patterns for the neutrino mass hierarchy are possible. $\Delta m_{31}^2 > 0$ is referred to as normal hierarchy (*NH*) with $m_1 < m_2 < m_3$ and $\Delta m_{31}^2 < 0$ is defined as inverted hierarchy (*IH*) with $m_3 < m_1 < m_2$. Combining Eq.(7) and Eq.(8), there is the following relation

$$2\nu_\Delta^2 \geq 1.96 \times 10^4 \left(\frac{200 GeV}{m_H}\right)^2 [V_{MNS} diag(0, \Delta m_{21}^2, \Delta m_{31}^2) V_{MNS}^+]_{e\mu}. \quad (9)$$

In our following numerical estimation, we will consider this constraint on the triplet *VEV* ν_Δ to calculate the cross sections for single production of the doubly charged *Higgs* boson H^{--} via $e\gamma$ collision at the *ILC* and *LHeC*.

3. Single production of the doubly charged Higgs boson H^{--} at the ILC

The *LHC* can generate very massive new particles and will essentially enlarge the possibilities of testing for new physics effects, while the *ILC* is also need to complement

the probe of the new particles with detailed measurement. For the *ILC*, the center of mass (*c.m.*) energy $\sqrt{s} = 0.5 - 1 \text{ TeV}$ and the typical integrated luminosity $\mathcal{L}_{int} = 0.5 - 1 \text{ ab}^{-1}$ is currently being designed [13, 14]. An unique feature of the *ILC* is that it can be transformed to $\gamma\gamma$ or $e\gamma$ collision with the photon beam generated by laser-scattering method. The effective luminosity and energy of the $\gamma\gamma$ and $e\gamma$ collisions are expected to be comparable to those of the *ILC*. In some scenarios, they are the best instrument for discovery of the new physics signatures. In this section, we will consider single production of the doubly charged *Higgs* boson H^{--} predicted by the *HTM* via $e\gamma$ collision.




Figure 1: The *Feynman* diagrams for the partonic process $e^-(p_1) + \gamma(p_2) \rightarrow l^+(p_3) + H^{--}(p_4)$ in the HTM.

From discussions given in section 2, we can see that the doubly charged *Higgs* boson H^{--} can be produced via $e\gamma$ collision associated with a positive lepton. The relevant *Feynman* diagrams are shown in Fig.1, in which l^+ denotes the lepton e^+, μ^+ or τ^+ . For the process $e^-(p_1) + \gamma(p_2) \rightarrow l^+(p_3) + H^{--}(p_4)$, the renormalization amplitude can be written as

$$\begin{aligned}
M &= M_a + M_b + M_c = -\frac{2eh_{el}}{(p_1 + p_2)^2} v^T(p_3) C^{-1} P_L (\not{p}_1 + \not{p}_2) \gamma^\mu u(p_1) \varepsilon_\mu(p_2) \\
&\quad - \frac{4eh_{el}}{(p_2 - p_4)^2 - m_H^2} v^T(p_3) C^{-1} P_L u(p_1) (2p_4 - p_2)^\mu \varepsilon_\mu(p_2) \\
&\quad - \frac{2eh_{el}}{(p_1 - p_4)^2} v^T(p_3) \gamma^\mu (\not{p}_1 - \not{p}_4) C^{-1} P_L u(p_1) \varepsilon_\mu(p_2).
\end{aligned} \tag{10}$$

Here $P_L = (1 - \gamma_5)/2$ is the right-handed projection operator and $\varepsilon_\mu(p_2)$ is the polarization vector of the photon γ . The explicit expressions for the *Yukawa* coupling constants h_{ee} , $h_{e\mu}$, and $h_{e\tau}$ can be written as [8, 9]

$$h_{ee} = \frac{1}{\sqrt{2}\nu_\Delta} [m_1 e^{-i\varphi_1} c_{12}^2 c_{13}^2 + m_2 s_{12}^2 c_{13}^2 + m_3 e^{i(2\delta - \varphi_2)} s_{13}^2], \tag{11}$$

$$h_{e\mu} = \frac{1}{\sqrt{2}\nu_\Delta} [m_1 e^{-i\varphi_1} c_{12} c_{13} (-s_{12} c_{23} - e^{-i\delta} c_{12} s_{13} s_{23}) + m_2 s_{12} c_{13} (c_{12} c_{23} - e^{-i\delta} s_{12} s_{13} s_{23}) + m_3 e^{i(\delta-\varphi_2)} s_{13} c_{13} s_{23}], \quad (12)$$

$$h_{e\tau} = \frac{1}{\sqrt{2}\nu_\Delta} [m_1 e^{-i\varphi_1} c_{12} c_{13} (s_{12} s_{23} - e^{-i\delta} c_{12} c_{23} s_{13}) + m_2 c_{13} s_{12} (-c_{12} s_{23} - e^{-i\delta} s_{12} s_{13} c_{23}) + m_3 e^{i(\delta-\varphi_2)} s_{13} c_{13} c_{23}]. \quad (13)$$




Figure 2: At the *ILC*, the production cross section σ_1 for the subprocess $e\gamma \rightarrow e^+ H^{--}$ as a function m_0 for different values of the m_H in the *NH*(a) and *IH*(b) cases.

After calculating the cross section $\hat{\sigma}(\hat{s})$ for the subprocess $e\gamma \rightarrow l^+ H^{--}$, the effective cross section $\sigma_1(s)$ at the *ILC* with the *c.m.* energy $\sqrt{s} = 1\text{TeV}$ can be obtained by folding the cross section $\hat{\sigma}(\hat{s})$ with the photon distribution function $f_{\gamma/e}(x)$ [21]

$$\sigma_1(s) = \int_{m_H^2/s}^{0.83} \hat{\sigma}(\hat{s}) f_{\gamma/e}(x) dx, \quad (14)$$

where $x = \hat{s}/s$, in which $\sqrt{\hat{s}}$ is the *c.m.* energy of the subprocess $e^-\gamma \rightarrow l^+ H^{--}$. In above equation, we have neglected the mass of the lepton l^+ due to $m_H \gg m_{l^+}$.

It is obvious that the production cross section σ_1 is dependent on the neutrino oscillation parameters, the triplet *VEV* ν_Δ , and the mass parameter m_H . Direct searches for $H^{\pm\pm}$ have been performed at *LEP* [22], *Tevatron* [23], and *HERA* [24] via the production mechanisms $e^+e^- \rightarrow H^{++}H^{--}$ and $q\bar{q} \rightarrow H^{++}H^{--}$. Both of these production




Figure 3: Same as Fig.2 but for the subprocess $e\gamma \rightarrow \mu^+ H^{--}$.

mechanisms are depend on only one unknown parameter m_H . The lower mass limits for m_H from the Tevatron searches are $m_H > 110\text{GeV} \sim 150\text{GeV}$ [23]. In this paper, we take m_H as a free parameter and assume that its value is in the range of $300\text{GeV} \sim 600\text{GeV}$. The current experimental upper limit for the LFV process $\mu \rightarrow e\gamma$ has give severe constraint on the triplet VEV ν_Δ , as shown in Eq.(9). In our numerical estimation, we take the minimal value given by Eq.(9). In this case, the production cross section σ_1 is only dependent on m_H and the neutrino oscillation parameters.

There are nine neutrino oscillation parameters: three neutrino masses m_i (or m_{ij}), three mixing angles θ_{ij} , the Dirac phase δ and two Majorana phases φ_1 and φ_2 . According the current constraints on the neutrinos and mixing parameters from neutrino oscillation experiments [25], we use the following values in this article

$$\begin{aligned}
\Delta m_{21}^2 &= m_2^2 - m_1^2 \simeq 7.59 \pm 0.20 \begin{pmatrix} +0.61 \\ -0.69 \end{pmatrix} \times 10^{-5} eV^2, \\
\Delta m_{31}^2 &= m_3^2 - m_1^2 = -2.36 \pm 0.11 (\pm 0.37) \times 10^{-3} eV^2 \text{ for } IH, \\
\Delta m_{31}^2 &= m_3^2 - m_1^2 = 2.46 \pm 0.12 (\pm 0.37) \times 10^{-3} eV^2 \text{ for } NH, \\
\theta_{12} &= 34.4 \pm 1.0 \begin{pmatrix} +3.2 \\ -2.9 \end{pmatrix}^\circ, \quad \theta_{23} = 42.8_{-2.9}^{+4.7} \begin{pmatrix} +10.7 \\ -7.3 \end{pmatrix}^\circ,
\end{aligned} \tag{15}$$

$$\theta_{13} = 5.6^{+3.0}_{-2.7} (\leq 12.5)^{\circ}. \quad (16)$$

Information on the mass m_0 of the lightest neutrino, the Dirac and Majorana phases can not be obtained from neutrino oscillation experiments. To simply our paper, we will take $\delta = 0$ and $\varphi_1 = \varphi_2 = 0$ to calculate the cross sections for single production of H^{--} in the cases of NH ($m_0 = m_1 < m_2 < m_3$) and IH ($m_0 = m_3 < m_1 < m_2$). For the free parameter m_0 , we will assume its value in the range of $0.001\text{eV} \leq m_0 \leq 1\text{eV}$.




Figure 4: Same as Fig.2 but for the subprocess $e\gamma \rightarrow \tau^+ H^{--}$.

From above discussions we can see that the production cross section for the process $e\gamma \rightarrow l^+ H^{--}$ depends on two free parameters m_0 and m_H . In our numerical estimation, we take the cross section σ_1 as a function of the lightest neutrino mass m_0 for three values of m_H . Our numerical results are summarized in Fig.2, Fig.3, and Fig.4, which correspond the subprocesses $e\gamma \rightarrow e^+ H^{--}$, $e\gamma \rightarrow \mu^+ H^{--}$, and $e\gamma \rightarrow \tau^+ H^{--}$, respectively. One can see from these figures that the value of the production cross section σ_1 increases as m_H increases, which is because the minimal value of the triplet VEV ν_Δ is proportional to the factor $1/m_H$. In both of the NH and IH cases, the production cross section $\sigma_1(\mu H)$ for the subprocess $e\gamma \rightarrow \mu^+ H^{--}$ is approximately equal to that for the subprocess $e\gamma \rightarrow \tau^+ H^{--}$ and their values are smaller than 16fb in most of the parameter space of the HTM . This is due to $h_{e\mu} \approx h_{e\tau}$, which reflects the neutrino mixing patterns. However, for the subprocess $e\gamma \rightarrow e^+ H^{--}$, it is not this case. For the NH spectrum, its cross section

$\sigma_1(eH)$ is larger or smaller than the cross section $\sigma_1(\mu H)$ (or $\sigma_1(\tau H)$), which depends on the mass m_0 of the lightest neutrino. For the IH spectrum, the cross section $\sigma_1(eH)$ is always larger than the cross section $\sigma_1(\mu H)$ (or $\sigma_1(\tau H)$). For $0.001\text{eV} \leq m_0 \leq 1\text{eV}$ and $300\text{GeV} \leq m_H \leq 600\text{GeV}$, the values of the cross sections for production of H^{--} associated with a positron e^+ via $e\gamma$ collision at the *ILC* with $\sqrt{s} = 1\text{TeV}$ are in the ranges of $1.7\text{fb} \sim 17.9\text{fb}$ and $34.8\text{fb} \sim 72.6\text{fb}$ corresponding to the *NH* and *IH* cases, respectively.




Figure 5: The number of the $\bar{e}\mu\mu$ (a) and $\bar{e}ee$ (b) events at the *ILC*.

Ref. [9] has shown that, for $\nu_\Delta < 0.1\text{MeV}$, the leptonic decays of $H^{\pm\pm}$ are the dominant decay modes, while the decay $H^{\pm\pm} \rightarrow W^\pm W^\pm$ is negligible. In the context of the *HTM*, the leptonic decay channels $H^{\pm\pm} \rightarrow l^\pm l^\pm$ have been extensively studied in Refs. [11, 26] and the direct connections between their branching ratios and the triplet *Yukawa* couplings h_{ij} (i.e. the neutrino oscillation parameters) are discussed in these papers. For the leptonic decays of $H^{\pm\pm}$, the possible decay modes are ee , $\mu\mu$, $\tau\tau$, $e\mu$, $e\tau$ and $\mu\tau$. In the case of $\delta = 0$ and $\varphi_1 = \varphi_2 = 0$, $\mu\mu$ and $\tau\tau$ are the dominate decay modes, their branching ratios are roughly equal to each other for the *NH* spectrum and $m_1 < 0.1\text{eV}$. While for the *IH* mass spectrum and $0.001\text{eV} \leq m_3 \leq 0.1\text{eV}$, the doubly charged *Higgs* boson H^{--} mainly decays to ee and its branching ratio is $32\% \leq Br(H^{--} \rightarrow ee) \leq 53\%$. Thus, for the final states generated by the subprocess $e\gamma \rightarrow e^+ H^{--}$, we consider the three

leptons $\bar{e}\mu\mu$ and $\bar{e}ee$ for the NH and IH cases, respectively. The number of the $\bar{e}\mu\mu$ and $\bar{e}ee$ events at the ILC with $\sqrt{s} = 1TeV$ and the integrated luminosity $\mathcal{L}_{int} = 500fb^{-1}$ are shown in Fig.5, where we have taken m_0 in the range of $0.001eV \sim 0.1eV$. From this figure, one can see that there will be several hundreds and up to ten thousands three lepton events to be generated per year in future ILC experiments. It is obvious that this is not the realistic number of the observed events. To take into account detector acceptance we should impose appropriate cuts. For example, the observed leptons must carry a minimum energy and respect a suitable rapidity cut, the angles of the observed leptons relative to the beam, θ_e, θ_μ , must be restricted to some ranges [27]. Detailed analysis is needed, which is beyond the scope of this paper.

It is well known that, at the ILC , the LFV processes can provide extremely clear signatures and are experimentally interesting. The leptonic decays of the doubly charged *Higgs* bosons give rise to same-sign lepton pairs, which can generate distinct experimental signals. The three lepton states $\bar{e}\mu\mu$ and $\bar{e}ee$ generated by the subprocess $e\gamma \rightarrow e^+H^{--}$ are almost free of the SM backgrounds at the ILC , which have been investigated in Ref. [27]. Thus, as long as the triplet VEV ν_Δ is enough small, the doubly charged *Higgs* boson H^{--} predicted by the HTM might be detected in future ILC experiments.

4. Single production of the doubly charged Higgs boson H^{--} at the LHeC

Recently, the high-energy ep collision has been considered at the LHC , which is called the $LHeC$ [15]. For the $LHeC$, the energy E_p of the incoming proton is given by the LHC beam, and the energy E_e of the incoming electron is in the range of $50 - 200GeV$, corresponding to the $c.m.$ energies of $\sqrt{s} = 2\sqrt{E_p E_e} \approx 1.18 - 2.37TeV$. Its anticipated integrated luminosity is at the order of $10 - 100fb^{-1}$ depending on the energy of the incoming electron and the design. Several studies have been performed to discuss the possibility of detecting the *Higgs* boson at the $LHeC$ [28]. In this section, we consider single production of the doubly charged *Higgs* boson H^{--} via $e\gamma$ collision at the $LHeC$.

It is well known that the equivalent photon approximation (EPA) can be successfully to describe most of the processes involving photon exchange [29]. A significant fraction of pp collision at the LHC will involve quasi-real photon interactions occurring at energies

well beyond the electroweak energy scale. The quasi-real photons emitted by the incoming protons have low virtuality Q^2 and scattered with small angles from the beam pipe. The photon spectrum depending on virtuality Q^2 and energy E_γ can be described by the following relation [30, 31]

$$dN = \frac{\alpha}{\pi} \frac{dE_\gamma}{E_\gamma} \frac{dQ^2}{Q^2} \left[\left(1 - \frac{E_\gamma}{E_p}\right) \left(1 - \frac{Q_{min}^2}{Q^2}\right) F_E + \frac{E_\gamma^2}{2E_p^2} F_M \right] \quad (17)$$

with

$$Q_{min}^2 = \frac{m_p^2 E_\gamma^2}{E_p(E_p - E_\gamma)}, \quad F_E = \frac{4m_p^2 G_E^2 + Q^2 G_M^2}{4m_p^2 + Q^2}, \quad (18)$$

$$G_E^2 = \frac{G_M^2}{\mu_p^2} = \left(1 + \frac{Q^2}{Q_0^2}\right)^4, \quad F_M = G_M^2, \quad Q_0^2 = 0.71 \text{ GeV}^2. \quad (19)$$

Where E_p is the energy of the incoming proton which is related to the photon energy by $E_\gamma = \xi E_p$ and m_p is the mass of the proton. Thus, ξ is the proton momentum fraction carried by the photon. The magnetic moment of the proton is $\mu_p^2 = 7.78$, F_E and F_M are functions of the electric and magnetic form factors. Electromagnetic form factors are steeply falling as a function of Q^2 . Then, the Q^2 integrated photon flux can be written as

$$f(E_\gamma) = \int_{Q_{min}^2}^{Q_{max}^2} \frac{dN}{dE_\gamma dQ^2} dQ^2, \quad (20)$$

where $Q_{max}^2 \approx 2 - 4 \text{ GeV}^2$. Since the contribution to the above integral formula is very small for $Q_{max}^2 > 2 \text{ GeV}^2$, in our numerical estimation, we will take $Q_{max}^2 \approx 2 \text{ GeV}^2$.

At the $LHeC$, the effective production cross section $\sigma_2(lH)$ for the subprocess $e\gamma \rightarrow l^+ H^{--}$ can be written as

$$\sigma_2(lH) = \int_{(m_H+m_l)^2/s}^{\xi_{max}} \sigma(\hat{s}) f(\xi E_p) d\xi, \quad (21)$$

where $\hat{s} = 4E_e E_\gamma = \xi s$ with $E_e = 140 \text{ GeV}$ and $E_p = 7 \text{ TeV}$. To detect the intact proton or its loss energy, the forward proton detectors are needed. The *ATLAS* forward physics (*AFP*) collaboration will have forward detectors with an acceptance of $0.0015 < \xi < 0.15$ [31]. The acceptance of the *CMS-TOTEM* forward detectors can reach $0.0015 < \xi < 0.5$ [32]. In our numerical estimation, we will take $\xi_{max} = 0.5$ and $\xi_{min} = (m_H + m_l)^2/s$, which corresponds the *c.m.* energy of the subprocess $e\gamma \rightarrow l^+ H^{--}$, $\sqrt{\hat{s}} \geq m_H + m_l$.




Figure 6: At the $LHeC$, the production cross section σ_2 for the subprocess $e\gamma \rightarrow e^+H^{--}$ as a function m_0 for different values of the m_H in the $NH(a)$ and $IH(b)$ cases.

Our numerical results are summarized in Fig.6 and Fig.7, which plot the production cross sections for the subprocesses $e\gamma \rightarrow e^+H^{--}$ and $e\gamma \rightarrow \mu^+H^{--}$ at the $LHeC$ as functions of the lightest neutrino mass m_0 for three values of the mass m_H of the doubly charged *Higgs* boson. In these figures we have shown our results for the NH and IH cases. Our calculation shows that, at the $LHeC$, the effective production cross section for the subprocess $e\gamma \rightarrow \mu^+H^{--}$ is roughly equal to that for the subprocess $e\gamma \rightarrow \tau^+H^{--}$, which is similar with the conclusion at the ILC . Thus, we have not given the contours for the subprocess $e\gamma \rightarrow \tau^+H^{--}$. From these figures, one can see that the production cross section of the subprocess $e\gamma \rightarrow l^+H^{--}$ at the $LHeC$ is smaller than that for the ILC . For $0.001eV \leq m_0 \leq 1eV$ and $300GeV \leq m_H \leq 600GeV$, the values of the production cross section $\sigma_2(eH)$ and $\sigma_2(\mu H)$ are in the ranges of $0.048fb \sim 0.51fb$ and $0.095fb \sim 0.43fb$, respectively, for the NH spectrum, while are in the ranges of $0.97fb \sim 2.03fb$ and $0.11fb \sim 0.45fb$, respectively, for the IH spectrum. If we assume the integrated luminosity $L_{int} = 100fb^{-1}$ of the $LHeC$ with $\sqrt{s} = 1.96TeV$, which corresponds $E_e = 140GeV$ and $E_p = 7TeV$, then there will be several and up to hundreds e^+H^{--} -events to be generated per year.

The process $ep \rightarrow e^+H^{--} + X$ with H^{--} decaying to ee can give rise to number




Figure 7: Same as Fig.6 but for the subprocess $e\gamma \rightarrow \mu^+ H^{--}$.

of signal events with same-sign electron pair and an isolated positron, which is almost free of the SM backgrounds. Thus, it is also possible to detect the signatures of the doubly charged *Higgs* boson H^{--} via its production associated with a positron at the *LHeC*. Certainly, detailed confirmation of the observability of the signals generated by the *HTM* at the *LHeC* experiments would require Monte-Carlo simulations of the signals and backgrounds, which is beyond the scope of this paper.

5. Conclusions and discussions

Doubly charged *Higgs* bosons ($H^{\pm\pm}$) appear in some popular new physics models beyond the SM , which can accommodate neutrino masses. This kind of new particles have distinct experimental signals through their decay to same-sign lepton pairs. Their observation in future high energy collider experiments would be a clear evidence of new physics. Thus, searching for $H^{\pm\pm}$ is one of the main goals of current and future high energy collider experiments.

The *HTM* is one of the attractive new physics models, in which a *Higgs* triplet is introduced and the neutrino masses can be obtained by the product of the triple VEV ν_Δ and the triplet *Yukawa* coupling h_{ij} . This model predicts the existence of the doubly charged *Higgs* bosons ($H^{\pm\pm}$). In this paper, we consider production of H^{--} associated with a positive lepton l^+ via $e\gamma$ collision at the *ILC* and *LHeC*. The production cross

sections are dependent on the neutrino oscillation parameters, the triplet VEV ν_Δ , and the mass parameter m_H . In our numerical estimations, we use the lower bound on ν_Δ^2 given by the LFV process $\mu \rightarrow e\gamma$, assume $\delta = 0$ and $\varphi_1 = \varphi_2 = 0$, and take the experimental measurement values for the parameters s_{ij} , c_{ij} , and m_{ij} . In this case, the production cross sections depend only on two free parameters, the mass parameter m_H and the lightest neutrino mass m_0 . From our numerical results, we can obtain following conclusions.

(1) The cross section for production of the doubly charged *Higgs* boson H^{--} associated with a positive lepton l^+ at the *ILC* is generally larger than that at the *LHeC*. The production cross sections for the *IH* mass spectrum are larger than those for the *NH* mass spectrum at these two kinds of collider experiments.

(2) Because of $h_{e\mu} \simeq h_{e\tau}$, the production cross section of the subprocess $e\gamma \rightarrow \mu^+ H^{--}$ is roughly equal to that of the subprocess $e\gamma \rightarrow \tau^+ H^{--}$ at both the *ILC* and *LHeC* experiments. For the *IH* mass spectrum, the cross section $\sigma(eH)$ is always larger than $\sigma(\mu H)$, while for the *NH* mass spectrum, $\sigma(eH)$ is larger or smaller than $\sigma(\mu H)$ which depends on the value of the lightest neutrino mass m_0 .

(3) For $0.001eV \leq m_0 \leq 1eV$ and $300GeV \leq m_H \leq 600GeV$, the values of the cross section $\sigma(eH)$ at the *ILC* with $\sqrt{s} = 1TeV$ are in the ranges of $1.7fb - 17.9fb$ and $34.8fb - 72.6fb$ corresponding with the *NH* and *IH* cases, respectively. While their values at the *LHeC* with $\sqrt{s} = 1.97TeV$ are in the ranges of $0.048fb - 0.51fb$ and $0.97fb - 2.03fb$ for the *NH* and *IH* cases, respectively.

(4) For the leptonic decays of H^{--} , the process $e\gamma \rightarrow l^+ H^{--}$ can produce three lepton final state, which is almost free of the *SM* backgrounds. In the context of the *HTM*, this process can only generate number of three lepton events at the *LHeC*, while, with reasonable values of the relevant free parameters, it can give rise to large number of three lepton events at the *ILC*. For instance, for $0.001eV \leq m_0 \leq 0.1eV$ and $300GeV \leq m_H \leq 600GeV$, the subprocess $e\gamma \rightarrow l^+ H^{--}$ with H^{--} decaying to ee can generate $8.0 \times 10^3 - 1.17 \times 10^4$ $\bar{e}ee$ events at the *ILC* experiment with $\sqrt{s} = 1000GeV$ and the integrated luminosity $\mathcal{L}_{int} = 500fb^{-1}$ in the *IH* mass spectrum. Thus, the possible

signals of H^{--} predicted the HTM might be detected via this process in the future ILC experiments.

In our numerical estimation, we have taken the minimal value of the triplet VEV ν_Δ demanded by the current upper experimental bound for the LFV process $\mu \rightarrow e\gamma$. However, the constraint on the HTM from the process $\mu \rightarrow ee\bar{e}$ is generally stronger than that from the process $\mu \rightarrow e\gamma$. For example, Ref. [12] has shown that, for $\varphi_1 = \varphi_2 = 0$ and $\sin^2 2\theta_{13} = 0$, the processes $\mu \rightarrow ee\bar{e}$ and $\mu \rightarrow e\gamma$ demand $\nu_\Delta m_H \geq 400 eV \cdot GeV$ and $\nu_\Delta m_H \geq 100 eV \cdot GeV$, respectively. It is obvious that the minimal value of ν_Δ demanded by $\mu \rightarrow e\gamma$ is smaller than that for $\mu \rightarrow ee\bar{e}$. Thus, if we use the constraints on the free parameters of the HTM coming from the process $\mu \rightarrow ee\bar{e}$, the production cross sections and the number of the signal events will be decreased at least by one order of magnitude. However, as long as $\nu_\Delta < 10^{-5} GeV$, there will be several $\bar{e}ee$ events to be generated at the ILC .

It has been shown [26] that the Majorana phases φ_1 and φ_2 have large effects on the branching ratios $Br(H^{\pm\pm} \rightarrow l_i^\pm l_j^\pm)$. Thus, φ_1 and φ_2 also have effects on the cross section for production of the doubly charged *Higgs* boson H^{--} associated with a positive lepton l^+ . If we vary the values of φ_1 and φ_2 , the numerical results for the cross sections $\sigma(eH)$, $\sigma(\mu H)$, and $\sigma(\tau H)$ are also changed. However, our physical conclusions are not changed.

Certainly, the doubly charged *Higgs* boson H^{++} can also be singly produced at the ILC and the $LHeC$ via the charge-conjugation processes of the corresponding processes for H^{--} . Similar to above calculation, we can give the values of the production cross sections for H^{++} . Thus, the conclusions for H^{--} also apply to H^{++} .

If the doubly charged *Higgs* bosons $H^{\pm\pm}$ are indeed found at the LHC , various consistency checks will have to be performed. One of the important tasks is to exactly calculate and precisely measure the branching ratios $Br(H^{\pm\pm} \rightarrow l_i^\pm l_j^\pm)$ and see whether their values are consistent with a neutrino mass matrix from oscillation data. We expect that our work will be helpful to test the Higgs triplet mechanism for neutrino masses and further to distinguish different new physics models.

Acknowledgments

This work is supported in part by the National Natural Science Foundation of China under Grant No.10975067, the Specialized Research Fund for the Doctoral Program of Higher Education(SRFDP) (No.200801650002).

References

- [1] C. K. Jung et al., *Ann. Rev. Nucl. Part. Sci.* **51**, 451(2001); Q. R. Ahmad et al.[SNO Collaboration], *Phys. Rev. Lett.* **89**, 011301(2002); K. Eguchi et al.[KamLAND Collaboration], *Phys. Rev. Lett.* **90**, 021802(2003); M. H. Ahn et al.[K2K Collaboration], *Phys. Rev. Lett.* **90**, 041801(2003).
- [2] V. Barger, D. Marfatia, and K. Whisnant, *Int. J. Mod. Phys. E***12**, 569(2003); M. C. Gonzalez-Garcia and M. Maltoni, *Phys. Rept.* **460**, 1(2008); R. N. Mohapatra and A. Y. Smirnov, *Ann. Rev. Nucl. Part. Sci.* **56**,569(2006); A. Strumia and F. Vissani, arXiv:hep-ph/0606054; Z. Z. Xing, *Int. J. Mod. Phys. A***23**, 4255(2008).
- [3] W. Konetschny and W. Kummer, *Phys. Lett. B***70**, 433(1977); T. P. Cheng and L. F. Li, *Phys. Rev. D***22**, 2860(1980); M. Magg and C. Wetterich, *Phys. Lett. B***94**, 61(1980).
- [4] J. Schechter and J. W. F. Valle, *Phys. Rev. D***22**, 2227(1980).
- [5] V. D. Barger, H. Baer, W. Y. Keung and R. J. N. Phillips, *Phys. Rev. D***26**, 218(1982).
- [6] J. F. Gunion, J. Grifols, A. Mendez, B. Kayser and F. I. Olness, *Phys. Rev. D***40**, 1546(1989); J. F. Gunion, C. Loomis and K. T. Pitts, *hep-ph/9610237*; K. Huitu, J. Maalampi, A. Pietila and M. Raidal, *Nucl. Phys. B***487**, 27(1997); J. Maalampi and N. Romanenko, *Phys. Lett. B***532**, 202(2002); M. Muhlleitner and M. Spira, *Phys. Rev. D***68**, 117701(2003); G. Azuelos, K. Benslama and J. Ferland, *J. Phys. G***32**,

- 73(2006); T. Rommerskirchen and T. Hebbeker, *J. Phys. G***34**, N47(2007); T. Han, B. Mukhopadhyaya, Z. Si and K. Wang, *Phys. Rev. D***76**, 075013(2007).
- [7] B. Dion, T. Gregoire, D. London, L. Marleau and H. Nadeau, *Phys. Rev. D***59**, 075006(1999); A. G. Akeroyd and M. Aoki, *Phys. Rev. D***72**, 035011(2005).
- [8] A. G. Akeroyd, M. Aoki, H. Sugiyama, *Phys. Rev. D***77**, 075010(2008); A. G. Akeroyd, Cheng-Wei Chiang, *Phys. Rev. D***80**, 113010(2009).
- [9] P. Fileviez Perez, T. Han, G. Y. Huang, T. Li and K. Wang, *Phys. Rev. D***78**, 015018(2008).
- [10] F. del Aguila and J. A. Aguilar-Saavedra, *Nucl. Phys. B***813**, 22(2009); A. G. Akeroyd, C.-W. Chiang, N. Gaur, *arXiv: 1009.2780* [hep-ph].
- [11] E. Ma, M. Raidal and U. Sarkar, *Phys. Rev. Lett.* **85**, 3769(2000); E. Ma, M. Raidal and U. Sarkar, *Nucl. Phys. B***615**, 313(2001); E. J. Chun, K. Y. Lee and S. C. Park, *Phys. Lett. B***566**, 142(2003); M. Kakizaki, Y. Ogura and F. Shima, *Phys. Lett. B***566**, 210(2003).
- [12] A. G. Akeroyd, M. Aoki, H. Sugiyama, *Phys. Rev. D***79**, 113010(2009); T. Fukuyama, H. Sugiyama, K. Tsumura, *JHEP* **1003**, 044(2010).
- [13] T. Abe et al.[American Linear Collider Group], *hep-ex/0106057*; J. A. Aguilar-Saavedra et al. [ECFA/DESY LC Physics Working Group],*hep-ph/0106315*; Koh Abe et al.[ACFA Linear Collider Working Group], *hep-ph/0109166*; ILC Technical Review Committee, second report, 2003, SLAC-R-606, February 2003.
- [14] G. Aarons et al. [ILC Collaboration], "International Linear Collider Reference Design Report Volume 2: Physics at the ILC," *arXiv: 0709.1893* [hep-ph]; Brau et al. [ILC Collaboration], "ILC Reference Design Report Volume 1 - Executive Summary," *arXiv: 0712.1950* [physics.acc-ph].

- [15] J. B. Dainton, M. Klein, P. Newman, E. Perez, and F. Willeke, *JINST* **1**, P10001(2006), *hep-ex/0603016*; P. Newman, *Nucl. Phys. Proc. Suppl.* **191**, 307(2009).
- [16] S. Chakrabarti, D. Choudhury, R. M. Godbole, B. Mukhopadhyaya, *Phys. Lett. B* **434**, 347(1998); E. M. Gregores, A. Gusso, S. F. Novaes, *Phys. Rev. D* **64**, 015004(2001); S. Godfrey, P. Kalyniak, N. Romanenko, *Phys. Rev. D* **65**, 033009(2002); Chong-Xing Yue, Shuang Zhao, Wei Ma, *Nucl. Phys. B* **784**, 36(2007).
- [17] Z. Maki, M. Nakagawa and S. Sakata, *Prog. Theor. Phys.* **28**, 870(1962).
- [18] S. M. Bilenky, J. Hosek and S. T. Petcov, *Phys. Lett. B* **94**, 495(1980); M. Doi, T. Kotani, H. Nishiura, K. Okuda and E. Takasugi, *Phys. Lett. B* **102**, 323(1981).
- [19] C. S. Lim and T. Inami, *Prog. Theor. Phys.* **67**, 1569(1982); A. Pich, A. Santamaria and J. Bernabeu, *Phys. Lett. B* **148**, 229(1984); M. L. Swartz, *Phys. Rev. D* **40**, 1521(1989); M. Raidal and A. Santamaria, *Phys. Lett. B* **421**, 250(1998).
- [20] M. L. Brooks et al. [MEGA Collaboration], *Phys. Rev. Lett.* **83**, 1521(1999).
- [21] F. Ginzburg *et al.*, *Nucl. Instrum* **219**, 5(1984); L. Ginzburg, G. Kotkin, H. Spiesberger, *Fortschr. Phys.* **34**, 687(1986); V. Telnov, *Nucl. Instrum. Methods Phys. Res. A* **294**, 72(1990).
- [22] G. Abbiendi et al. [OPAL Collaboration], *Phys. Lett. B* **526**, 221(2002); P. Achard et al. [L3 Collaboration], *Phys. Lett. B* **576**, 18(2003); J. Abdallah et al. [DELPHI Collaboration], *Phys. Lett. B* **552**, 127(2003).
- [23] V. M. Abazov et al. [D0 Collaboration], *Phys. Rev. Lett.* **93**, 141801(2004); D. E. Acosta et al. [CDF Collaboration], *Phys. Rev. Lett.* **93**, 221802(2004); V. M. Abazov et al. [D0 Collaboration], *Phys. Rev. Lett.* **101**, 071803(2008); T. Aaltonen et al. [The CDF Collaboration], *Phys. Rev. Lett.* **101**, 121801(2008).
- [24] A. Aktas et al. [H1 Collaboration], *Phys. Lett. B* **638**, 432(2006).

- [25] M. C. Gonzalez-Garcia, M. Maltoni, J. Salvado, *JHEP* **1004**, 056(2010).
- [26] J. Garayoa and T. Schwetz, *JHEP* **0803**, 009(2008); A. G. Akeroyd, M. Aoki and H. Sugiyama, *Phys. Rev. D* **77**, 075010(2008); M. Kadastik, M. Raidal and L. Rebane, *Phys. Rev. D* **77**, 115023(2008).
- [27] S. Chakrabarti, D. Choudhury, R. M. Godbole, B. Mukhopadhyaya, *Phys. Lett. B* **434**, 347(1998); E. M. Gregores, A. Gusso, S. F. Novaes, *Phys. Rev. D* **64**, 015004(2001); S. Godfrey, P. Kalyniak, N. Romanenko, *Phys. Rev. D* **65**, 033009(2002); M. Frank, K. Huitu, S. K. Rai, *Phys. Rev. D* **77**, 015006(2008).
- [28] U. Klein, Talk given at the 2nd CERN-ECFA-NuPECC Workshop on the LHeC, Divonnes, France, September 2009; M. Kuze, Talk given at the 2nd CERN-ECFA-NuPECC Workshop on the LHeC, Divonnes, France, September 2009; T. Han, B. Mellado, *Phys. Rev. D* **82**, 016009(2010); B. Jager, *Phys. Rev. D* **81**, 054018(2010).
- [29] V. M. Budnev, I. F. Ginzburg, G.V. Meledin and V. G. Serbo, *Phys. Rep.* **15**, 181(1975).
- [30] G. Baur, K. Hencken, D. Trautmann, S. Sadowsky and Y. Kharlov, *Phys. Rep.* **364**, 359(2002).
- [31] M. G. Albrow et al. (FP420 R and D Collaboration), *arXiv*: **0806.0302** [hep-ph].
- [32] V. Avati and K. Osterberg, Report No. CERN-TOTEM-NOTE-2005-002, 2006; O. Kepka and C. Royon, *Phys. Rev. D* **78**, 073005(2008).